

On the Rationality of Algebraic Tori of Norm Type

Shizuo Endo

Teikyo University Hachioji Tokyo 192-0352 Japan

metadata, citation and similar papers at core.ac.uk

Received February 20, 1999

1

Let G be a finite group. A G -module is called a *permutation module* if it is isomorphic to $\bigoplus_{i=1}^r \mathbf{Z}G/G_i$, where G_i , $1 \leq i \leq r$, are subgroups of G . A G -module M is called a *quasi-permutation module* if there exists an exact sequence

$$0 \rightarrow M \rightarrow S \rightarrow S' \rightarrow 0,$$

where S and S' are permutation G -modules. The dual module $\text{Hom}_{\mathbf{Z}}(M, \mathbf{Z})$ of a G -module M is denoted by M° .

Let k be a field and let F be an extension field of k . Then F is said to be *rational* over k if it is generated by a finite number of elements of F which are algebraically independent over k and to be *stably rational* over k if there exists an extension field of F which is rational over each of k and F . Further, F is said to be *retract rational* over k if it is the quotient field of an integral domain $A \supseteq k$ such that there are k -algebra homomorphisms

$$\begin{aligned} \phi : A &\rightarrow k[X_1, X_2, \dots, X_n][1/s] \quad \text{and} \\ \psi : k[X_1, X_2, \dots, X_n][1/s] &\rightarrow A, \end{aligned}$$

where $k[X_1, X_2, \dots, X_n][1/s]$ is the localization of a polynomial ring $k[X_1, X_2, \dots, X_n]$ with variables X_1, X_2, \dots, X_n over k with respect to $0 \neq s \in k[X_1, X_2, \dots, X_n]$ and $\psi \cdot \phi$ is the identity on A . It is easy to see that *rational* \Rightarrow *stably rational* \Rightarrow *retract rational*.

Let G be a finite group and let M be a G -module with a \mathbf{Z} -free basis u_1, u_2, \dots, u_n . Let k be a field and let K be a Galois extension of k with group G . Define the action of G on the rational function field

$K(X_1, X_2, \dots, X_n)$ with variables X_1, X_2, \dots, X_n over K , as an extension of the action of G on K , as follows. For each $\sigma \in G$,

$$\sigma(X_i) = \prod_{j=1}^n X_j^{m_{ij}}, \quad 1 \leq i \leq n,$$

when $\sigma u_i = \sum_{j=1}^n m_{ij} u_j$, $m_{ij} \in \mathbf{Z}$, and denote $K(X_1, X_2, \dots, X_n)$ with this action of G by $K(M)$.

As is well known, there is an algebraic torus T defined over k and split over K such that the character group of T is isomorphic to M as G -modules, and the invariant subfield $K(M)^G$ of $K(M)$ can be identified with the function field of T over k .

We now have the following fundamental

PROPOSITION. *Let G be a finite group and let k be a field. Let K be a Galois extension of k with group G and let M be a \mathbf{Z} -free and \mathbf{Z} -finitely generated G -module (i.e., a G -lattice). Then:*

- (1) (e.g., [2, 1.6]) *M is a quasi-permutation G -module if and only if $K(M)^G$ is stably rational over k .*
- (2) [7, 3.14] *M is a direct summand of a quasi-permutation G -module if and only if $K(M)^G$ is retract rational over k .*

Let p be a prime, and let P be an elementary abelian p -group of order p^m , $m \geq 1$. Let P_i , $1 \leq i \leq r$, be distinct subgroups of index p in P , and let $\varepsilon_i: \mathbf{Z}P/P_i \rightarrow \mathbf{Z}$ be the augmentation maps. Further, for h_1, h_2, \dots, h_r , ≥ 1 , let

$$\Phi = (\varepsilon_1^{h_1}, \varepsilon_2^{h_2}, \dots, \varepsilon_r^{h_r}): \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i} \rightarrow \mathbf{Z}$$

and set $L = \text{Ker } \Phi$.

The main result of this note is the following

THEOREM 1. (1) *In case of $p = 2$, L° is a (direct summand of a) quasi-permutation P -module if and only if $r = 1, 2$.*

(2) *In case of $p \neq 2$, L° is a (direct summand of a) quasi-permutation P -module if and only if $r = 1$.*

By the Proposition, Theorem 1 can be restated as follows:

THEOREM 2. *Let G , P , and L be as above and let K be a Galois extension of a field k with group P . Then:*

- (1) *In case of $p = 2$, $K(L^\circ)^G$ is stably (retract) rational over k if and only if $r = 1, 2$.*

(2) In case of $p \neq 2$, $K(L^\circ)^G$ is stably (retract) rational over k if and only if $r = 1$.

It is noted that the algebraic torus corresponding to L° defined over k and split over K is of *norm type*.

Part (1) of Theorem 2 is an answer to the question asked the late professor T. Miyata by A. Merkurjev in 1982 [6]. It should be noted that Theorem 1 was obtained in 1982 and reported without proof in [4].

2

We now begin to prove Theorem 1. In order to prove this, we need to consider a more general situation. Let P , P_i , and ε_i , $1 \leq i \leq r$, be as above, and define the homomorphism $\delta: \mathbf{Z} \rightarrow \mathbf{Z}$ by $\delta(1) = p$. For $h_1, h_2, \dots, h_r \geq 1$ and $h \geq 0$, let

$$\Phi = (\varepsilon_1^{h_1}, \varepsilon_2^{h_2}, \dots, \varepsilon_r^{h_r}, \delta): \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i} \oplus \mathbf{Z}^h \rightarrow \mathbf{Z},$$

and set $L = \text{Ker } \Phi$. Further, let

$$\Phi' = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r): \bigoplus_{i=1}^r \mathbf{Z}P/P_i \rightarrow \mathbf{Z},$$

and set $L' = \text{Ker } \Phi'$. Then the key lemma is given as follows:

LEMMA. $L \cong L' \oplus \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i-1} \oplus \mathbf{Z}^h$.

Proof. First, define the homomorphism $\nu_i: \mathbf{Z}P/P_i \rightarrow (\mathbf{Z}P/P_i)^{h_i}$ by $\nu_i(a_i) = (a_i, 0, 0, \dots, 0)$ for each $1 \leq i \leq r$, and set

$$\nu = (\nu_1, \nu_2, \dots, \nu_r): \bigoplus_{i=1}^r \mathbf{Z}P/P_i \rightarrow \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i} \subseteq \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i} \oplus \mathbf{Z}^h.$$

Next, define the homomorphism $\mu_i: (\mathbf{Z}P/P_i)^{h_i} \rightarrow \mathbf{Z}P/P_i$ by $\mu_i((a_{i1}, a_{i2}, \dots, a_{ih_i})) = \sum_{j=1}^{h_i} a_{ij}$ for each $1 \leq i \leq r$ and the homomorphism $\eta: \mathbf{Z}^h \rightarrow \bigoplus_{i=1}^r \mathbf{Z}P/P_i$ by $\eta((b_1, b_2, \dots, b_h)) = ((\sum_{i=1}^h b_i s, 0, 0, \dots, 0))$ ($\eta = 0$ when $h = 0$), where s denotes the sum of all elements of P/P_1 in $\mathbf{Z}P/P_1$, and set

$$\mu = (\mu_1, \mu_2, \dots, \mu_r, \eta): \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i} \oplus \mathbf{Z}^h \rightarrow \bigoplus_{i=1}^r \mathbf{Z}P/P_i.$$

Then $\mu \cdot \nu$ is the identity on $\bigoplus_{i=1}^r \mathbf{Z}P/P_i$ and it can be easily seen that $\nu(L') \subseteq L$ and $\mu(L) \subseteq L'$. Therefore, denoting the restrictions of ν , μ to L' , L by $\tilde{\nu}$, $\tilde{\mu}$, respectively, we get homomorphisms $\tilde{\nu}: L' \rightarrow L$ and

$\tilde{\mu}: L \rightarrow L'$ such that $\tilde{\mu} \cdot \tilde{\nu}$ is the identity on L' . This shows that $L \cong L' \oplus \text{Coker } \tilde{\nu}$. However, as is easily seen, $\text{Coker } \tilde{\nu} \cong \text{Coker } \nu \cong \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i-1} \oplus \mathbf{Z}^h$. Thus this concludes that $L \cong L' \oplus \bigoplus_{i=1}^r (\mathbf{Z}P/P_i)^{h_i-1} \oplus \mathbf{Z}^h$.

By this lemma we have only to consider the case where $h_1 = h_2 = \cdots = h_r = 1$, that is, the exact sequence

$$0 \rightarrow L \rightarrow \bigoplus_{i=1}^r \mathbf{Z}P/P_i \rightarrow \mathbf{Z} \rightarrow 0, \quad (*)$$

where P_1, P_2, \dots, P_r are distinct subgroups of index p in P .

It is known that L° is a quasi-permutation P -module for each of the cases (a) $r = 1$ and (b) $r = 2$ and $p = 2$ [6]. However, for completeness, we give a proof of it.

Assume first that $r = 1$. Then P/P_1 is cyclic, and so $L^\circ \cong L$. Hence L° is clearly a quasi-permutation P -module. Next, assume that $r = 2$ and $p = 2$. Then $P/P_1 \cap P_2$ is the Klein four group, and therefore, regarding $P/P_1 \cap P_2$ as P , we can set $P = \langle \sigma, \tau \mid \sigma^2 = \tau^2 = 1, \sigma\tau = \tau\sigma \rangle$ and $P_1 = \langle \sigma \rangle$, $P_2 = \langle \tau \rangle$. Now we have $\text{Ker } \Phi = \mathbf{Z}P(1, -1)$. Define the epimorphism $\Psi = (\mu, \nu): \mathbf{Z}P \oplus \mathbf{Z} \rightarrow L$ by $\mu(1) = (1, -1)$ and $\nu(1) = (1 + \tau, 1 + \sigma)$. Then, as is easily seen, $\text{Ker } \Psi = \{((a + b\sigma)(1 + \sigma\tau), -(a + b)) \mid a, b \in \mathbf{Z}\}$, and so $\text{Ker } \Psi \cong \mathbf{Z}P/\langle \sigma\tau \rangle$. Thus we have the exact sequence $0 \rightarrow \mathbf{Z}P/\langle \sigma\tau \rangle \rightarrow \mathbf{Z}P \oplus \mathbf{Z} \rightarrow L \rightarrow 0$, which shows that L° is a quasi-permutation P -module.

3

Consider here the case here $p = 2$ and $r \geq 3$. This case can be divided into the following:

- (1) $[P : P_{i_1} \cap P_{i_2} \cap P_{i_3}] = 4$ for some $i_1 < i_2 < i_3$.
- (2) $[P : P_{i_1} \cap P_{i_2} \cap P_{i_3}] = 8$ for any $i_1 < i_2 < i_3$.

Set $P' = P_{i_1} \cap P_{i_2} \cap P_{i_3}$ in both cases and restrict the exact sequence $(*)$ to P' . Then we obtain the exact sequence

$$0 \rightarrow L^{P'} \rightarrow \mathbf{Z}P/P_{i_1} \oplus \mathbf{Z}P/P_{i_2} \oplus \mathbf{Z}P/P_{i_3} \oplus \bigoplus_{i \neq i_1, i_2, i_3} (\mathbf{Z}P/P_i)^{P'} \rightarrow \mathbf{Z} \rightarrow 0.$$

It is noted that L° is not (a direct summand of) a quasi-permutation P -module if $(L^{P'})^\circ$ is not (a direct summand of) a quasi-permutation P/P' -module. Regarding P/P' as P and applying the lemma to the above

exact sequence, the cases (1) and (2) reduce, respectively, to the following

- (i) $P = \langle \sigma, \tau \mid \sigma^2 = \tau^2 = 1, \sigma\tau = \tau\sigma \rangle$, the Klein four group,

$$0 \rightarrow L \rightarrow \mathbf{Z}P/\langle \sigma \rangle \oplus \mathbf{Z}P/\langle \tau \rangle \oplus \mathbf{Z}P/\langle \sigma\tau \rangle \rightarrow \mathbf{Z} \rightarrow 0.$$

- (ii) $P = \langle \rho, \sigma, \tau \mid \rho^2 = \sigma^2 = \tau^2 = 1, \rho\sigma = \sigma\rho, \sigma\tau = \tau\sigma, \tau\rho = \rho\tau \rangle$

(a) $0 \rightarrow L \rightarrow \mathbf{Z}P/\langle \rho, \sigma \rangle \oplus \mathbf{Z}P/\langle \tau, \rho \rangle \oplus \mathbf{Z}P/\langle \sigma, \tau \rangle \rightarrow \mathbf{Z} \rightarrow 0.$

(b) $0 \rightarrow L \rightarrow \mathbf{Z}P/\langle \rho, \sigma \rangle \oplus \mathbf{Z}P/\langle \tau, \rho \rangle \oplus \mathbf{Z}P/\langle \sigma, \tau \rangle \oplus \mathbf{Z}P/\langle \rho\sigma, \rho\tau \rangle \rightarrow \mathbf{Z} \rightarrow 0.$

We have only to show for these cases that L° is not a direct summand of any quasi-permutation P -module. For brevity, a direct summand of a quasi-permutation module is called a *d-quasi-permutation* module.

It should be noted that the case (i) was examined by Miyata (unpublished, 1974) and Hürlimann [5]. We give here a proof of this case for completeness. Let P be as in the case (i) and let I be the augmentation ideal of $\mathbf{Z}P$. Define the homomorphism $\phi: \mathbf{Z}P/\langle \sigma \rangle \oplus \mathbf{Z}P/\langle \tau \rangle \oplus \mathbf{Z}P/\langle \sigma\tau \rangle \rightarrow \mathbf{Z}P$ as

$$\phi((1, 0, 0)) = 1 + \sigma, \quad \phi((0, 1, 0)) = 1 + \tau, \quad \phi((0, 0, 1)) = 1 + \sigma\tau.$$

Restricting ϕ to L , we obtain the epimorphism $\tilde{\phi}: L \rightarrow I$. It is easy to see that $\text{Ker } \tilde{\phi} \cong \mathbf{Z}^2$ is a trivial P -module. Hence we have the exact sequence

$$0 \rightarrow \mathbf{Z}^2 \rightarrow L \rightarrow I \rightarrow 0.$$

However, it is known [1, 3] that I° is not a d-quasi-permutation module. This concludes that L° is not a d-quasi-permutation module.

Next, consider the case (ii), (a). We restrict the exact sequence to the subgroup $P' = \langle \rho\sigma, \rho\tau \rangle$ of P . Then we see that $\mathbf{Z}P/\langle \rho, \sigma \rangle \oplus \mathbf{Z}P/\langle \rho, \tau \rangle \oplus \mathbf{Z}P/\langle \sigma, \tau \rangle \cong \mathbf{Z}P'/\langle \rho\sigma \rangle \oplus \mathbf{Z}P'/\langle \rho\tau \rangle \oplus \mathbf{Z}P'/\langle \sigma\tau \rangle$ as P' -modules. This shows by the case (i) that L° is not a d-quasi-permutation P' -module. Thus L° is also not a d-quasi-permutation P -module.

The case (ii), (b) is slightly complicated. We set $M = \mathbf{Z}P/\langle \rho \rangle \oplus \mathbf{Z}P/\langle \sigma \rangle \oplus \mathbf{Z}P/\langle \tau \rangle \oplus \mathbf{Z}P/\langle \rho\sigma \rangle \oplus \mathbf{Z}P/\langle \sigma\tau \rangle \oplus \mathbf{Z}P/\langle \tau\rho \rangle \oplus \mathbf{Z}^3$ and $N = \mathbf{Z}P/\langle \rho, \sigma \rangle \oplus \mathbf{Z}P/\langle \tau, \rho \rangle \oplus \mathbf{Z}P/\langle \sigma, \tau \rangle \oplus \mathbf{Z}P/\langle \rho\sigma, \rho\tau \rangle$. Define the 9 homomorphisms as

$$\mathbf{Z}P/\langle \rho \rangle \rightarrow N \quad \text{by } 1 \rightarrow (1, -1, 0, 0)$$

$$\mathbf{Z}P/\langle \sigma \rangle \rightarrow N \quad \text{by } 1 \rightarrow (1, 0, -1, 0)$$

$$\begin{array}{ll}
\mathbf{Z}P/\langle \tau \rangle \rightarrow N & \text{by } 1 \rightarrow (0, 1, -1, 0) \\
\mathbf{Z}P/\langle \rho\sigma \rangle \rightarrow N & \text{by } 1 \rightarrow (1, 0, 0, -1) \\
\mathbf{Z}P/\langle \sigma\tau \rangle \rightarrow N & \text{by } 1 \rightarrow (0, 0, 1, -1) \\
\mathbf{Z}P/\langle \tau\rho \rangle \rightarrow N & \text{by } 1 \rightarrow (0, 1, 0, -1) \\
\mathbf{Z} \rightarrow N & \text{by } 1 \rightarrow (1 + \tau, -(1 + \sigma), 0, 0) \\
\mathbf{Z} \rightarrow N & \text{by } 1 \rightarrow (1 + \tau, 0, -(1 + \rho), 0) \\
\mathbf{Z} \rightarrow N & \text{by } 1 \rightarrow (1 + \tau, 0, 0, -(1 + \sigma\tau)).
\end{array}$$

By adding these homomorphisms, we obtain the homomorphism $\Psi: M \rightarrow N$. It is easy to see that the image of Ψ coincides with L . Set $H = \text{Ker } \Psi$. Then we have the exact sequence

$$0 \rightarrow H \rightarrow M \rightarrow L \rightarrow 0.$$

By direct computation we show that $M^{P'} \rightarrow L^{P'}$ is surjective for each subgroup P' of P . Therefore it follows that $H^1(P', H) = 0$ for each subgroup P' of P , where $H^i(\ , \)$ denotes the Tate cohomology group.

Assume that L° is a d-quasi-permutation. Then, by (the proof of) [3, 1.6], there is an exact sequence

$$0 \rightarrow T \rightarrow D \rightarrow L \rightarrow 0,$$

where T is a permutation module and D is a direct summand of permutation module. Taking the pullback of $M \rightarrow L \leftarrow D$, we obtain $H \oplus D \cong T \oplus M$. Since P is a 2-group, H is locally isomorphic to a permutation module S . We denote the completion of a module V at 2 by V^* . Then we have $H^* \cong S^*$. Set $\lambda = \rho\sigma\tau$.

Considering $\langle \lambda \rangle$ -invariants we have $(L^*)^{\langle \lambda \rangle} \cong (\mathbf{Z}^*)^3$ and

$$(H^*)^{\langle \lambda \rangle} \cong (\mathbf{Z}^*P/\langle \rho\sigma, \tau \rangle \oplus \mathbf{Z}^*P/\langle \tau\rho, \sigma \rangle \oplus \mathbf{Z}^*P/\langle \sigma\tau, \rho \rangle)^2.$$

Since $\text{rank}_Z H = 20$ and $\text{rank}_Z H^P = 6$, we have

$$H^* \cong \mathbf{Z}^*P/P_1 \oplus \mathbf{Z}^*P/P_2 \oplus \mathbf{Z}^*P/P_3 \oplus \mathbf{Z}^*P/P_4 \oplus \mathbf{Z}^*P/P'_1 \oplus \mathbf{Z}^*P/P'_2,$$

where $|P_1| = |P_2| = |P_3| = |P_4| = 2$ and $|P'_1| = |P'_2| = 4$. Using the cohomology sequence of the exact sequence $0 \rightarrow L \rightarrow N \rightarrow \mathbf{Z} \rightarrow 0$, we see that $H^1(P, L) \cong \mathbf{Z}/2\mathbf{Z}$ and $H^0(P, L) \cong (\mathbf{Z}/4\mathbf{Z})^3$. Further, taking the cohomology sequence of the exact sequence $0 \rightarrow H \rightarrow M \rightarrow L \rightarrow 0$, we have the exact sequence $H^0(P, H) \rightarrow H^0(P, M) \rightarrow H^0(P, L) \rightarrow H^1(P, H) = 0$. By direct computation we also have $H^0(P, H) \cong (\mathbf{Z}/2\mathbf{Z})^4 \oplus (\mathbf{Z}/4\mathbf{Z})^2$ and $H^0(P, M) \cong (\mathbf{Z}/2\mathbf{Z})^4 \oplus (\mathbf{Z}/8\mathbf{Z})^3$. Therefore we obtain the exact sequence

$$(\mathbf{Z}/2\mathbf{Z})^4 \oplus (\mathbf{Z}/4\mathbf{Z})^2 \rightarrow (\mathbf{Z}/2\mathbf{Z})^6 \oplus (\mathbf{Z}/8\mathbf{Z})^3 \rightarrow (\mathbf{Z}/4\mathbf{Z})^3 \rightarrow 0.$$

However, it is easy to see that the kernel of $(\mathbf{Z}/2\mathbf{Z})^6 \oplus (\mathbf{Z}/8\mathbf{Z})^3 \rightarrow (\mathbf{Z}/4\mathbf{Z})^3$

is isomorphic to $(\mathbf{Z}/2\mathbf{Z})^9$. This is a contradiction, because $(\mathbf{Z}/2\mathbf{Z})^4 \oplus (\mathbf{Z}/4\mathbf{Z})^2$ never maps onto $(\mathbf{Z}/2\mathbf{Z})^9$. Thus L° is not a d-quasi-permutation.

4

Finally consider the case where $p \neq 2$ and $r \geq 2$. We now prove that L° is not a d-quasi-permutation. Restricting the exact sequence $(*)$ to the subgroup $P' = P_1 \cap P_2$, regarding P/P' as P and using the lemma as in the case where $p = 2$, we can reduce this case to the one where $|P| = p^2$. From now on, assume that P is of order p^2 . Under this assumption it is noted that there exist exactly $p + 1$ subgroups of order p in P . Let P_i , $1 \leq i \leq r$, be distinct subgroups of order p in P and set $N = \bigoplus_{i=1}^r \mathbf{Z}P/P_i$. Let $\varepsilon_i : \mathbf{Z}P/P_i \rightarrow \mathbf{Z}$, $1 \leq i \leq r$, be the augmentation maps and let

$$\Phi = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r) : N \rightarrow \mathbf{Z}.$$

Set $L = \text{Ker } \Phi$. Then we have the exact sequence

$$0 \rightarrow L \rightarrow N \rightarrow \mathbf{Z} \rightarrow 0.$$

For $1 \leq i \leq r$, set $P/P_i = \langle \rho_i \rangle$ and $s_i = \sum_{j=0}^{p-1} \rho_i^j \in \mathbf{Z}P/P_i$. Define the homomorphisms as

$$\begin{array}{ll} \mathbf{Z}P \rightarrow N & \text{by } 1 \rightarrow (1, -1, 0, \dots, 0) \\ \mathbf{Z}P \rightarrow N & \text{by } 1 \rightarrow (1, 0, -1, 0, \dots, 0) \\ & \dots \\ & \dots \\ \mathbf{Z}P \rightarrow N & \text{by } 1 \rightarrow (1, 0, \dots, 0, -1) \\ \mathbf{Z} \rightarrow N & \text{by } 1 \rightarrow (s_1, -s_2, 0, \dots, 0) \\ \mathbf{Z} \rightarrow N & \text{by } 1 \rightarrow (s_1, 0, -s_3, 0, \dots, 0) \\ & \dots \\ & \dots \\ \mathbf{Z} \rightarrow N & \text{by } 1 \rightarrow (s_1, 0, 0, \dots, 0, -s_r) \\ \mathbf{Z}P/P_1 \rightarrow N & \text{by } 1 \rightarrow (\rho_1 - 1, 0, \dots, 0) \\ \mathbf{Z}P/P_2 \rightarrow N & \text{by } 1 \rightarrow (0, \rho_2 - 1, 0, \dots, 0) \\ & \dots \\ & \dots \\ \mathbf{Z}P/P_r \rightarrow N & \text{by } 1 \rightarrow (0, 0, \dots, 0, \rho_r - 1). \end{array}$$

Set $M = \mathbf{Z}P^{r-1} \oplus \mathbf{Z}^{r-1} \oplus \bigoplus_{i=1}^r \mathbf{Z}P/P_i$ and define the homomorphism $\Psi : M \rightarrow N$ as the sum of the above homomorphisms. Then it is easy to

see that $\text{Im } \Psi = L$. Further set $H = \text{Ker } \Psi$. Then we have the exact sequence

$$0 \rightarrow H \rightarrow M \rightarrow L \rightarrow 0.$$

As is easily seen, we have

$$\begin{aligned} \text{rank}_Z N &= pr, & \text{rank}_Z N^P &= r, \\ \text{rank}_Z L &= pr - 1, & \text{rank}_Z L^P &= r - 1, \\ \text{rank}_Z M &= (p^2 + 1)(r - 1) + pr, & \text{rank}_Z M^P &= 3r - 2, \\ \text{rank}_Z H &= p^2(r - 1) + r, & \text{rank}_Z H^P &= 2r - 1. \end{aligned}$$

By direct computation we can show that $M^{P'} \rightarrow L^{P'}$ is surjective for each subgroup P' of P . Hence it follows that $H^1(P', H) = 0$ for each subgroup P' of P .

Assume that L° is a d-quasi-permutation. Then, as in the proof of the case where $p = 2$, it follows that H is a direct summand of a permutation module. Since P is a p -group, H is locally isomorphic to a permutation module S .

We denote the completion of a module V at p by V^* . Then we can decompose $H^* \cong S^* \cong (\mathbf{Z}^*P)^s \oplus (S')^*$, where S' does not have $\mathbf{Z}P$ as a direct summand. It is easy to see that $\text{rank}_Z S' = p^2(r - s - 1) + r$ and $\text{rank}_Z (S')^P = 2r - s - 1$. Since each indecomposable component of S' is of rank 1 or p over \mathbf{Z} , we have $\text{rank}_Z S' \leq p(2r - s - 1)$. If $r - s - 3 \geq 0$, it follows from the fact that $r \leq p + 1 < 2p$ that $\text{rank}_Z S' > p(2r - s - 1)$, which is a contradiction. Hence we have $r - s - 2 \leq 0$, that is, $r - 2 \leq s$. Since $s \leq r - 1$, this shows that $s = r - 2$ or $r - 1$.

Forming the pushout of $(S')^* \leftarrow S^* \rightarrow M^*$, we obtain the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & (\mathbf{Z}^*P)^s & = & (\mathbf{Z}^*P)^s & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & S^* & \longrightarrow & M^* & \longrightarrow & L^* \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & (S')^* & \longrightarrow & (M')^* & \longrightarrow & L^* \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0. & & \end{array}$$

Then the columns are split, and so $(M')^* = (\mathbf{Z}^*P)^{r-s-1} \oplus (\mathbf{Z}^*)^{r-1} \oplus \bigoplus_{i=1}^r \mathbf{Z}^*P/P_i$. Set $U^* = \sum_p (M^*)^{P'}$ where P' runs over all subgroups of order p in P . Since $(\mathbf{Z}^*)^{r-1} \oplus \bigoplus_{i=1}^r \mathbf{Z}^*P/P_i \subseteq U^*$, there exists an epimorphism $(\mathbf{Z}^*P)^{r-s-1} \rightarrow M^*/U^* \rightarrow L^*/\Psi^*(U^*)$. It can be shown that $\Psi^*(U^*) = \mathbf{Z}^*P(\rho_1 - 1, 0, \dots, 0) + \mathbf{Z}^*P(0, \rho_2 - 1, 0, \dots, 0) + \dots + \mathbf{Z}^*P(0, \dots, 0, \rho_r - 1) + \mathbf{Z}^*P(p, -p, 0, \dots, 0) + \mathbf{Z}^*P(p, 0, -p, 0, \dots, 0) + \dots + \mathbf{Z}^*P(p, 0, \dots, 0, -p)$. Hence it follows that $L^*/\Psi^*(U^*) \cong \mathbf{Z}/p\mathbf{Z}(1, -1, 0, \dots, 0) + \mathbf{Z}/p\mathbf{Z}(1, 0, -1, 0, \dots, 0) + \dots + \mathbf{Z}/p\mathbf{Z}(1, 0, \dots, 0, -1) \cong (\mathbf{Z}/p\mathbf{Z})^{r-1}$. Therefore we obtain an epimorphism $(\mathbf{Z}^*P)^{r-s-1} \rightarrow (\mathbf{Z}/p\mathbf{Z})^{r-1}$. However, $s = r - 2$ or $r - 1$, as seen above. This shows that $r = 2$ and $s = 0$. Then we have $\text{rank}_{\mathbf{Z}} H = p^2 + 2$ and $\text{rank}_{\mathbf{Z}} H^P = 3$. This is a contradiction, because $H^*(\cong (S')^*)$ does not have \mathbf{Z}^*P as a direct summand. Thus the proof is complete.

REFERENCES

1. C. Chevalley, On algebraic group varieties, *J. Math. Soc. Japan* **6** (1954), 303–324.
2. S. Endo and T. Miyata, Invariants of finite abelian groups, *J. Math. Soc. Japan* **25** (1973), 7–26.
3. S. Endo and T. Miyata, On a classification of the function fields of algebraic tori, *Nagoya Math. J.* **56** (1975), 85–104; Corrigenda, **79** (1980), 187–190.
4. S. Endo, Non-rationality of algebraic tori of norm type and its application to generic division rings, in “Proc. 22nd Symp. Ring Theory, Okayama Univ., 1989,” pp. 59–64.
5. W. Hürlimann, On algebraic tori of norm type, *Comment. Math. Helv.* **59** (1984), 539–549.
6. A. Merkurjev, letter to T. Miyata, October 25, 1982.
7. D. J. Saltman, Retract rational fields and cyclic Galois extensions, *Israel J. Math.* **47** (1984), 165–215.
8. V. E. Voskresenskiĭ, Algebraic groups and their birational invariants, in *Transl. Math. Monogr.*, Vol. 179, Amer. Math. Soc., Providence, 1998.